

## Some Topics on Weighted Generalized Inverse and Kronecker Product of Matrices

Zeyad Abdel Aziz Al Zhour & Adem Kilicman

*Department of Mathematics*

*Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu.*

*E-mail: zeyad1968@yahoo.com , akilicman@umt.edu.my*

### ABSTRACT

In this paper, we present two general representations for the weighted generalized inverse  $A_{d,w}$ , which extends earlier results on the Drazin inverse  $A_d$ , group inverse  $A_g$  and usual inverse  $A^{-1}$ . The first one concerns with the matrix expression involving Moore-Penrose inverse  $A^+$ . The second one holds on the Kronecker products of two and several matrices. Furthermore, some necessary and sufficient conditions for Drazin and weighted Drazin inverses are given for the reverse order law  $(AB)_d = B_d A_d$  and  $(AB)_{d,z} = B_{d,R} A_{d,w}$  to hold. Finally, we apply our result to present the solution of restricted singular matrix equations.

**Keywords:** Kronecker Product, Weighted Drazin Inverses, General algebraic structures, Index, Nilpotent matrix.

(2000 Mathematics Subject Classification: 15A69 ; 15A09.)

### INTRODUCTION AND PRELIMINARY RESULT

One of the important types of generalized inverses of matrices is the weighted Drazin inverse, which has several important applications such as, applications in singular differential, difference equations, Markov chains, statistical problems, control system analysis, curve fitting, numerical analysis and Kronecker product systems [e.g., 4, 7, 9, 13, 15, 16, 17]. Here we use the following notations. Let  $M_{m,n}$  be the set of all matrices over the complex number field  $\mathbb{C}$  and when  $m=n$ , we write  $M_n$  instead of  $M_{n,n}$ . For matrix  $A \in M_{m,n}$ , let  $A^*$  be the conjugate transpose of  $A$  and  $\text{rank}(A)$  be the rank of  $A$ . If  $A \in M_m$  is a given matrix, then the smallest non-negative integer  $k$  such that

$$\text{rank}(A^{k+1}) = \text{rank}(A^k) \tag{1}$$

is called the *index of  $A$*  and is denoted by  $\text{Ind}(A)=k$ .

It is well known that the Drazin inverse (DI) of  $A \in M_m$  with  $Ind(A)=k$  is defined to be the unique solution  $X \in M_m$  satisfying the following three matrix equations:

$$A^k X A = A^k, \quad X A X = X, \quad A X = X A \tag{2}$$

and is often denoted by  $X = A_d$ . Note that the first equation in (2) can be written as  $A^{k+1} X = A^k$ . In particular, when  $Ind(A)=1$ , the Drazin inverse of  $A \in M_m$  is called the *group inverse* of  $A$ , and is often denoted by  $A_g$ , but when  $Ind(A)=0$  and  $A \in M_m$  is a non-singular matrix, then  $A_d = A^{-1}$ .

Wang [13] gave that for  $A \in M_m$  with  $Ind(A) = k$ ,

$$A^k (A_d)^k A^k = A^k, \quad (A_d)^k A^k (A_d)^k = (A_d)^k, \quad A^k (A_d)^k = (A_d)^k A^k \tag{3}$$

By the uniqueness of the DI, we have

$$(A^k)_d = (A_d)^k \tag{4}$$

For more properties concerning Drazin inverses, see [e.g., 3, 4, 10, 14].

Cline and Greville [5] extended the Drazin inverse of square matrix to rectangular matrix and called it as the weighted Drazin inverse (WDI). The WDI of  $A \in M_{m,n}$  with respect to the matrix  $W \in M_{n,m}$  is defined to be the unique solution  $X \in M_{m,n}$  of the following three matrix equations:

$$(AW)^{k+1} X W = (AW)^k, \quad X W A W X = X, \quad A W X = X W A \tag{5}$$

where

$$k = \max \{Ind(AW), Ind(WA)\} \tag{6}$$

and is often denoted by  $X = A_{d,W}$ . In particular, when  $A \in M_m$  and  $W = I_m$ , then  $A_{d,W}$  reduce to  $A_d$ , i.e.,  $A_d = A_{d,I_m}$ . If  $A \in M_m$  is non-singular square matrix and  $W = I_m$ , it is easily seen that  $Ind(A) = 0$  and  $A_{d,W} = A_d = A^{-1}$  satisfies the matrix equations (5).

The properties of WDI can be found in [e.g., 8, 18, 19]. Some notable properties are: If  $A \in M_{m,n}$  with respect to the matrix  $W \in M_{n,m}$  and  $k = \max \{Ind(AW), Ind(WA)\}$ , then:

$$i. \quad A_{d,W} = A \{(WA)_d\}^2 = \{(AW)_d\}^2 A \tag{7}$$

$$\text{ii} \quad A_{d,w}W = (AW)_d, \quad WA_{d,w} = (WA)_d \quad (8)$$

$$\text{iii} \quad WAWA_{d,w} = WA(WA)_d, \quad A_{d,w}WAW = (WA)_dAW \quad (9)$$

iv One closed-form solution of  $A_{d,w}$  for a rectangular matrix  $A \in M_{m,n}$

$$A_{d,w} = \begin{cases} \lim_{\alpha \rightarrow 0} (AW)^{l+2} + \alpha^2 I)^{-1} (AW)^l A & \text{if } l \geq k \\ \lim_{\alpha \rightarrow 0} A(WA)^l ((WA)^{l+2} + \alpha^2 I)^{-1} & \text{if } l \geq k \end{cases} \quad (10)$$

The *Moore-Penrose inverse* (MPI) is a generalization of the inverse of non-singular matrix to the inverse of a singular and rectangular matrix. The MPI of a matrix  $A \in M_{m,n}$  is defined to be the unique solution  $X \in M_{n,m}$  of the following four Penrose equations:

$$AXA = A, \quad XAX = A, \quad (AX)^* = AX, \quad (XA)^* = XA \quad (11)$$

and is often denoted by  $X = A^+$ .

Note that if  $A \in M_m$  is non-singular matrix, then  $A^+ = A^{-1}$ . Regarding various basic properties concerning MPI, see [e.g., 2, 3, 4, 10].

The *general algebraic structures* (GAS) of the matrices  $A \in M_{m,n}$ ,  $W \in M_{n,m}$ ,  $A^+$ ,  $W^+$ , and  $A_{d,w} \in M_{n,m}$  with  $k = \max\{Ind(AW), Ind(WA)\}$  are (see [e.g., 4,19,20,21]):

$$A = L \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} Q^{-1}, \quad W = Q \begin{bmatrix} W_{11} & 0 \\ 0 & W_{22} \end{bmatrix} L^{-1}, \quad A^+ = Q \begin{bmatrix} A_{11}^{-1} & 0 \\ 0 & 0 \end{bmatrix} L^{-1}, \quad (12)$$

$$W^+ = L \begin{bmatrix} W_{11}^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1}, \quad A_{d,w} = L \begin{bmatrix} (W_{11}A_{11}W_{11})^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1} \quad (13)$$

where  $L, Q, A_{11}, W_{11}$  are non-singular matrices, and  $A_{22}, W_{22}, A_{22}W_{22}, W_{22}A_{22}$  are nilpotent matrices (A matrix  $A \in M_n$  is called nilpotent if  $A^k = 0$  for some positive integer  $k$ ).

In particular, when  $A \in M_n$  with  $Ind(A) = k$ ,  $W = I_m$  and  $L = Q$ , then we have

$$A = L \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} L^{-1}, \quad A_d = \begin{bmatrix} A_{11}^{-1} & 0 \\ 0 & 0 \end{bmatrix} L^{-1} \tag{14}$$

where  $L$  and  $A_{11}$  are non-singular matrices, and  $A_{22}$  is a nilpotent matrix.

Greville [6] first studied the reverse order law for the Drazin inverse of the product of two matrices  $A$  and  $B \in M_n$ . He proved that  $(AB)_d = B_d A_d$  holds under the condition  $AB = BA$ .

Tian [12] gave a necessary and sufficient condition for the reverse order law  $(AB)_d = B_d A_d$  by using a rank identity. A similar result for reverse order law for Drazin inverse of general multiple matrix product was presented by Wang [14] as follows: Let  $A$  and  $B \in M_n$  be given with  $k = \max \{Ind(A), Ind(B), Inx(AB)\}$ . Then

$$(AB)_d = B_d A_d \tag{15}$$

if and only if

$$\text{rank} \begin{bmatrix} (-1)^n (AB)^{2k+1} & 0 & 0 & (AB)^k \\ 0 & 0 & A^{2k+1} & A^k \\ 0 & B^{2k+1} & B^k A^k & 0 \\ (AB)^k & B^k & 0 & 0 \end{bmatrix} = \text{rank}(A^k) + \text{rank}(B^k) + \text{rank}((AB)^k) \tag{16}$$

Finally, the Kronecker product of  $A = [a_{ij}] \in M_{m,n}$  and  $B = [b_{kl}] \in M_{p,q}$  is given by

$$A \otimes B = \left[ a_{ij} B \right]_{ij} \in M_{mp,nq} \tag{17}$$

where  $a_{ij} B \in M_{p,q}$  is the  $ij$ -th block.

For any compatible matrices  $A, B, C$  and  $D$ ; and any real number  $r$ , we shall make frequent use of the following properties of the Kronecker product (see [e.g.,1,2,7,11,22]):

- i. If  $AC$  and  $BD$  are well defined, then.

$$(A \otimes B)(C \otimes D) = AC \otimes BD \quad (18)$$

ii. If  $A$  and  $B$  are square positive (semi) definite matrices, then

$$(A \otimes B)^r = A^r \otimes B^r \quad (19)$$

iii.  $\text{rank}(A \otimes B) = \text{rank}(A)\text{rank}(B)$  (20)

iv.  $\text{Vec}(AXB^T) = (B \otimes A) \text{Vec}(X)$  (21)

where

$$\text{Vec}(X) = \begin{bmatrix} x_{11} & \dots & x_{m1} & x_{12} & \dots & x_{m2} & \dots & x_{1m} & \dots & x_{mm} \end{bmatrix}^T \quad (22)$$

denotes vectorization by columns of arbitrary matrix  $X \in M_{m,n}$

v. If  $A$  and  $B$  are nilpotent matrices, then  $A \otimes B$  is nilpotent matrix

vi. If  $A$  and  $B$  are unitary matrices, then  $A \otimes B$  is unitary matrix.

In this paper, some new matrix expressions involving the three kinds of generalized inverses of the Kronecker products matrices are established. In addition, by using the general algebraic structures of matrices (GAS), the necessary and sufficient conditions for Drazin and weighted Drazin inverses are also given for the reverse order laws  $(AB)_d = B_d A_d$  and  $(AB)_{d,z} = B_{d,k} A_{d,w}$  to hold. Finally, we apply our result to present the solution of restricted singular matrix equations (WAW)  $X(RBR)^T = C$ .

### MAIN RESULT

Observe that, in general, if  $A$  and  $B \in M_n$  are nilpotent matrices, then  $AB$  need not be nilpotent. As an example, let

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

It is easy to verify that  $A$  and  $B$  are nilpotent matrices, but  $AB = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  is not nilpotent, because  $(AB)^k = AB \neq 0$  for all positive integer  $k$ . This observation is

important to give a nilpotent condition when we use the GAS of matrices under usual product as follows:

*Theorem 1* : Let

$$A = L \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} L^{-1}, \quad B = L \begin{bmatrix} B_{11} & 0 \\ 0 & B_{22} \end{bmatrix} L^{-1}, \quad W = L \begin{bmatrix} W_{11} & 0 \\ 0 & W_{22} \end{bmatrix} L^{-1}, \quad (23)$$

$$R = L \begin{bmatrix} R_{11} & 0 \\ 0 & R_{22} \end{bmatrix} L^{-1}, \quad Z = L \begin{bmatrix} Z_{11} & 0 \\ 0 & Z_{22} \end{bmatrix} L^{-1} \quad (24)$$

be the general algebraic structures, respectively, of  $A, B, W, R$  and  $Z \in M_n$  with  $k = \max \{Ind(AW), Ind(WA), Ind(BR), Ind(RB), Ind(ABZ), Ind(ZAB)\}$ . Then

$$(AB)_{d,Z} = B_{d,R} A_{d,W} \quad (25)$$

if and only if  $A_{22} B_{22}$  is a nilpotent matrix and

$$(R_{11} B_{11} R_{11})^{-1} (W_{11} A_{11} W_{11})^{-1} = (Z_{11} A_{11} B_{11} Z_{11})^{-1} \quad (26)$$

Or equivalently

$$R_{11}^{-1} B_{11}^{-1} R_{11}^{-1} W_{11}^{-1} A_{11}^{-1} W_{11}^{-1} = Z_{11}^{-1} B_{11}^{-1} A_{11}^{-1} Z_{11}^{-1} \quad (27)$$

**Proof** : The GAS of  $A, B, W, R$  and  $Z$  in the assumptions assure that  $A_{11}, B_{11}, W_{11}, R_{11}, Z_{11}$  and  $L$  are non-singular matrices, and  $A_{22}, B_{22}, W_{22}, R_{22}$  and  $Z_{22}$  are nilpotent. Then it is well known that the GAS of  $A_{d,W}$  and  $B_{d,R}$  are given by

$$A_{d,W} = L \begin{bmatrix} (W_{11} A_{11} W_{11})^{-1} & 0 \\ 0 & 0 \end{bmatrix} L^{-1}, \quad B_{d,R} = L \begin{bmatrix} (R_{11} B_{11} R_{11})^{-1} & 0 \\ 0 & 0 \end{bmatrix} L^{-1} \quad (28)$$

Computation shows that

$$B_{d,R} A_{d,W} = L \begin{bmatrix} (R_{11} B_{11} R_{11})^{-1} (W_{11} A_{11} W_{11})^{-1} & 0 \\ 0 & 0 \end{bmatrix} L^{-1} \quad (29)$$

and

$$(AB)_{d,Z} = \left( L \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix} L^{-1} \right)_{d,Z} = L \begin{bmatrix} (Z_{11}A_{11}B_{11}Z_{11})^{-1} & 0 \\ 0 & 0 \end{bmatrix} L^{-1} \quad (30)$$

if and only if  $A_{22}B_{22}$  is a nilpotent matrix. It is clear from (29) and (30) that

$$(AB)_{d,Z} = B_{d,K}A_{d,W}$$

if and only if  $A_{22}B_{22}$  is a nilpotent matrix and

$$\left( \begin{bmatrix} R_{11} & B_{11} & R_{11} \end{bmatrix} \right)^{-1} \left( \begin{bmatrix} W_{11} & A_{11} & W_{11} \end{bmatrix} \right)^{-1} = \left( \begin{bmatrix} Z_{11} & A_{11} & B_{11} & Z_{11} \end{bmatrix} \right)^{-1}$$

This completes the proof of Theorem 1. ■

If we set  $W = R = Z = I_n$  in Theorem 1, we obtain the sufficient and necessary condition for the reverse order of Drazin inverse as follows.

*Corollary 1* Let

$$A = L \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} L^{-1}, B = L \begin{bmatrix} B_{11} & 0 \\ 0 & B_{22} \end{bmatrix} L^{-1}$$

be the GAS of  $A$  and  $B \in M_n$ , respectively, with  $k = \max \{Ind(A), Ind(B), Ind(AB)\}$ .

Then

$$(AB)_d = B_d A_d$$

if and only if  $A_{22}B_{22}$  is a nilpotent matrix.

Now, we can also apply the GAS in order to find a new representation of WDI as follows:

*Theorem 2:* Let  $A \in M_{m,n}$  and  $W \in M_{n,m}$  such that  $A_{22}W_{22}$  and  $W_{22}A_{22}$  are nilpotent matrices of index  $k$  in GAS form. Then the WDI of  $A$  with respect to the matrix  $W$  can be written as matrix expression involving MPI by

$$A_{d,W} = \left\{ (AW)^k \left[ (AW)^{2k+1} \right]^+ (AW)^k \right\} W^+ \quad (31)$$

where  $k = \max \{ind(AW), Ind(WA)\}$ .

**Proof:** Due to the GAS of  $A, A^+W, W^+$  and  $A_{d,W}$  there exists non-singular matrices  $L, A_{11}$  and  $W_{11}$ , and nilpotent matrices  $A_{22}$  and  $W_{22}$  such that

$$A = L \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} Q^{-1}, W = Q \begin{bmatrix} W_{11} & 0 \\ 0 & W_{22} \end{bmatrix} L^{-1}, W^+ = Q \begin{bmatrix} W_0^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1},$$

Since  $A_{22}W_{22}$  and  $W_{22}A_{22}$  are nilpotent matrices of index  $k$ , then  $(A_{22}W_{22})^k = 0$ , and it is easy to show that

$$(AW)^k = L \begin{bmatrix} (A_{11}W_{11})^k & 0 \\ 0 & 0 \end{bmatrix} L^{-1}, \quad [(AW)^{2k+1}]^+ = L \begin{bmatrix} (A_{11}W_{11})^{-2k-1} & 0 \\ 0 & 0 \end{bmatrix}$$

Computation shows that

$$\begin{aligned} & \left\{ (AW)^k [(AW)^{2k+1}]^+ (AW)^k \right\} W^+ \\ &= L \begin{bmatrix} (A_{11}W_{11})^k & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} (A_{11}W_{11})^{-2k-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} (A_{11}W_{11})^k & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} W_{11}^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1} \\ &= L \begin{bmatrix} (A_{11}W_{11})^k (A_{11}W_{11})^{-2k-1} (A_{11}W_{11})^k W_{11}^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1} \\ &= L \begin{bmatrix} (A_{11}W_{11})^{-1} W_{11}^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1} \\ &= L \begin{bmatrix} (W_{11}A_{11}W_{11})^{-1} & 0 \\ 0 & 0 \end{bmatrix} Q^{-1} \\ &= A_{d,w} \end{aligned}$$

This completes the proof of Theorem 2. ■

If  $A$  is a square matrix with  $Ind(A) = k$  and set  $W = I_n$  in Theorem 2, we obtain the following corollary which is given by Wang [14]:

Corollary 2 Let  $A \in M_n$  with  $Ind(A) = k$ , then

$$A_d = A^k \left( A^{2k+1} \right)^+ A^k \tag{32}$$

Theorem 3 Let  $A \in M_{m,n}$ ,  $W \in M_{n,m}$ ,  $B \in M_{p,q}$  and  $R \in M_{q,p}$  be matrices with

$$k_1 = \max \{ Ind(AW), Ind(WA) \}, k_2 = \max \{ Ind(BR), Ind(RB) \}$$

Also, let  $Z = W \otimes R$  and  $k = \max \{ k_1, k_2 \}$ . Then



$$\text{i. } \text{Ind} \{(A \otimes B)Z\} = k \quad (33)$$

$$\text{ii. } (A \otimes B)_{d,z} = A_{d,w} \otimes B_{d,R} \quad (34)$$

**Proof:** i. By assumptions, we have

$$\text{rank}(AW)^{k_1} = \text{rank}(AW)^{k_1+1}, \text{rank}(BR)^{k_2} = \text{rank}(BR)^{k_2+1}$$

From properties of Kronecker products, we have

$$\begin{aligned} \text{rank}\{(A \otimes B)Z\}^r &= \text{rank}\{(A \otimes B)(W \otimes R)\}^r = \text{rank}\{AW \otimes BR\}^r \\ &= \text{rank}\{AW\}^r \text{rank}\{BR\}^r . \end{aligned}$$

Similarly,

$$\text{rank}\{(A \otimes B)Z\}^{r+1} = \text{rank}\{AW\}^{r+1} \text{rank}\{BR\}^{r+1} .$$

It is obvious that the smallest non-negative integer such that

$$\text{rank}\{(A \otimes B)Z\}^{r+1} = \text{rank}\{(A \otimes B)Z\}^r .$$

is  $k = \max\{k_1, k_2\}$ . Hence (33) is true.

ii. Let  $X = A_{d,w} \otimes B_{d,R}$  and  $Z = W \otimes R$ . From properties of the Kronecker product and (5) we have

$$\begin{aligned} ((A \otimes B)Z)^{k+1} XZ &= ((A \otimes B)(W \otimes R))^{k+1} (A_{d,w} \otimes B_{d,R})(W \otimes R) \\ &= ((AW)^{k+1} A_{d,w} W) \otimes ((BR)^{k+1} B_{d,R} R) \\ &= (AW)^k \otimes (BR)^k = (AW \otimes BR)^k = ((A \otimes B)(W \otimes R))^k \\ &= \{(A \otimes B)Z\}^k \end{aligned} \quad (35)$$

$$\begin{aligned} XZ(A \otimes B)ZX &= (A_{d,w} \otimes B_{d,R})(W \otimes R)(A \otimes B)(W \otimes R)(A_{d,w} \otimes B_{d,R}) \\ &= (A_{d,w} W A W A_{d,w}) \otimes (B_{d,R} R B R B_{d,R}) = A_{d,w} \otimes B_{d,R} \\ &= X. \end{aligned} \quad (36)$$

$$\begin{aligned} (A \otimes B)ZX &= (A \otimes B)(W \otimes R)(A_{d,w} \otimes B_{d,R}) \\ &= A W A_{d,w} \otimes B R B_{d,R} = A_{d,w} W A \otimes B_{d,R} R B \\ &= (A_{d,w} \otimes B_{d,R})(W \otimes R)(A \otimes B) \\ &= XZ(A \otimes B) \end{aligned} \quad (37)$$

From (35)-(37) we can obtain (34) immediately. ■

If  $A$  and  $B$  are square matrices with  $Ind(A) = k_1$  and  $Ind(B) = k_2$ , respectively, and set  $W = I_m$  and  $R = I_n$  in Theorem 3 we obtain the following corollary which is given by Wang [13]:

*Corollary: 3* Let  $A \in M_m$  and  $B \in M_n$  with  $Ind(A) = k_1$  and  $Ind(B) = k_2$ , respectively.

Then

$$Ind\{(A \otimes B)\} = \max\{k_1, k_2\} \tag{38}$$

and

$$(A \otimes B)_d = A_d \otimes B_d \tag{39}$$

More particularly, if  $Ind(A) = Ind(B) = 1$ , then we have

$$(A \otimes B)_g = A_g \otimes B_g \tag{40}$$

*Corollary: 4* Let  $A_i \in M_{m(i),n(i)}$  and  $W_i \in M_{n(i),m(i)}$  ( $1 \leq i \leq r, r \geq 2$ ) be matrices with

$$k_i = \max\{Ind(A_i W_i), Ind(W_i A_i)\}, i = 1, 2, \dots, r.$$

Then

$$Ind\left\{\left(\prod_{i=1}^r A_i\right)Z\right\} = k, \tag{41}$$

and 
$$\left(\prod_{i=1}^r A_i\right)_{d,z} = \prod_{i=1}^r (A_i)_{d,w_i} \tag{42}$$

where  $k = \max\{k_1, k_2, \dots, k_r\}$  and  $Z = \prod_{i=1}^r W_i$ . In particular,

i. if  $A_i \in M_{m(i)}$  and  $W_i = I_{n_i}$  ( $1 \leq i \leq k, k \geq 2$ ), we then have

$$Ind\left(\prod_{i=1}^r A_i\right) = k \tag{43}$$

and

$$\left(\prod_{i=1}^r A_i\right)_d = \prod_{i=1}^r (A_i)_d \tag{44}$$

where  $k = \max\{Ind(A_i), i = 1, 2, \dots, r\}$

ii. if  $Ind(A_1) = Ind(A_2) = \dots = Ind(A_r) = 1$ , we then have

$$\left(\prod_{i=1}^r A_i\right)_g = \prod_{i=1}^r (A_i)_g \tag{45}$$

**Proof**

The proof of (41) is by induction on  $r$ . The base case (when  $r = 2$ ) has been established in (33) of Theorem 3. In the induction hypothesis, we assume that

$$\begin{aligned} \text{Ind} \left\{ \left( \prod_{i=1}^{r-1} A_i \right) \left( \prod_{i=1}^{r-1} W_i \right) \right\} &= \text{Ind} \left\{ \prod_{i=1}^{r-1} A_i W_i \right\} \\ &= \gamma = \max \{ k_1, k_2, \dots, k_{r-1} \} \end{aligned}$$

Now

$$\begin{aligned} \text{Ind} \left\{ \left( \prod_{i=1}^r A_i \right) Z \right\} &= \text{Ind} \left\{ \left( \prod_{i=1}^r A_i \right) \left( \prod_{i=1}^r W_i \right) \right\} \\ &= \text{Ind} \left\{ \prod_{i=1}^r A_i W_i \right\} \\ &= \text{Ind} \left\{ \left( \prod_{i=1}^{r-1} A_i W_i \right) \otimes (A_r W_r) \right\} \\ &= \max \{ \gamma, k_r \} = \max \{ k_1, k_2, \dots, k_r \} = k \end{aligned}$$

The proof of (42) is also by induction on  $r$ . The base case (when  $r=2$ ) has been established in (34) of Theorem 3. In the induction hypothesis, we assume that

$$\left( \prod_{i=1}^{r-1} A_i \right)_{d, \prod_{i=1}^{r-1} e_i W_i} = \prod_{i=1}^{r-1} \otimes (A_i)_{d, W_i}$$

Now

$$\begin{aligned} \left( \prod_{i=1}^r A_i \right)_{d, Z} &= \left[ \left( \prod_{i=1}^{r-1} A_i \right)_{d, \prod_{i=1}^{r-1} e_i W_i} \right] \otimes (A_r)_{d, W_r} \\ &= \left( \prod_{i=1}^{r-1} \otimes (A_i)_{d, W_i} \right) \otimes (A_r)_{d, W_r} \\ &= \prod_{i=1}^r \otimes (A_i)_{d, W_i} \end{aligned}$$

The proof of three special cases in (43)-(45) are straightforward. ■

One of the important application of Theorem 3 is that the weighted Drazin inverse of Kronecker product arise naturally in solving the so-called restricted singular matrix equations (RSME) as follows.

*Theorem 4:* Let  $A \in M_{m,n}, W \in M_{n,m}, B \in M_{p,q}, R \in M_{q,p}$  and  $C \in M_{n,q}$  be given constant matrices and  $X \in M_{m,p}$  be an unknown matrix to be solved. Also, let

$$L = R \otimes W, k_1 = \text{Ind}((B \otimes A)L), k_2 = \text{Ind}(L(B \otimes A)) \quad (46)$$

such that

$$r((B \otimes A)L)^{k_1} = r(L(B \otimes A))^{k_2}, \text{Vec}C \in R(L(B \otimes A))^{k_2}, \text{Vec}X \in R((B \otimes A)L)^{k_1} \quad (47)$$

Then the unique solution of the following *RSME*

$$(WAW)X(RBR)^T = C \quad (48)$$

is given by

$$X = A_{d,W}CB_{d,R}^T \quad (49)$$

**Proof**

Using identity (21) it is not difficult to transform (48) into the vector form as:

$$(L(B \otimes A)L)\text{Vec}X = \text{Vec}C \quad (50)$$

It is easy to verify under conditions (47) that the unique solution of (50) is

$$\begin{aligned} \text{Vec}X &= (B \otimes A)_{d,L} \text{Vec}C = (B_{d,R} \otimes A_{d,W})\text{Vec}C \\ &= \text{Vec}(A_{d,W}CB_{d,R}^T) \end{aligned}$$

which is the required result. ■

An important particular case of Theorem 4 is that when  $m = n, p = q, W = I_m$  and  $R = I_p$ , we obtain the following corollary:

*Corollary 5* : Let  $A \in M_m, B \in M_p$  and  $C \in M_{m,p}$  be given constant matrices and  $X \in M_{m,p}$  be an unknown matrix to be solved. Then the unique solution of the following *RSME*

$$AXB^T = C : \text{Vec}C, \text{Vec}X \in R(B \otimes A)^k, k = \text{Ind}(B \otimes A) \quad (51)$$

is given by

$$X = A_dCB_d^T \quad (52)$$

**CONCLUSION**

In this paper, we have presented two general representations for weighted Drazin inverse related to Moore-Penrose inverse and Kronecker product of two and several matrices. These representations are viewed as a generalization of Wang's results in [13, Lemma 1.1, and 14, Theorem 2.2]. Furthermore, some necessary and sufficient conditions for Drazin and weighted Drazin inverses are given for the reverse order law  $(AB)_d = B_dA_d$  and  $(AB)_{d,zm} = B_{d,R}A_{d,w}$  to hold. Although the results are applied to solve the restricted singular matrix equations, the idea adopted can be easily extended to

solve the coupled restricted singular matrix equations. It is natural to ask if we can extend our results to the Minkowski inverse in Minkowski space. This will be part of future research.

## REFERENCES

- AL ZHOUR, Z. and A. KILICMAN. 2006. Matrix equalities and inequalities involving Khatri-Rao and Tracy-Singh Sums. *J. of Ineq. in Pure and Appl. Math.* **7(1)** Article 34.
- AL ZHOUR, Z. and A. KILICMAN. 2006. Extensions and generalization inequalities involving the Khatri-Rao product of several positive matrices. *J. of Ineq. and Appl.* **2006** Article ID 80878: 1-20.
- BEN - ISRAEL, A. and T. N. E. GREVILLE. 1974. *Generalized Inverses: Theory and Applications*. New York: Wiley.
- CAMPBELL, S.L. and C.D. MEYER. 1999. *Generalized Inverses of Linear Transformations*. New York: Dover Publ. Inc.
- CLINE, R.E. and T.N.E. GREVILLE. 1980. A Drazin inverse for rectangular matrices. *Lin. Alg. and its Appl.* **29**: 53-62.
- GREVILLE, T.N.E. 1966. Note on generalized inverse of matrix product. *SIAM Rev.* **8**: 518-521.
- KILICMAN, A. and Z. AL ZHOUR. 2005. The general common exact solutions of coupled linear matrix and matrix differential equations. *J. Anal. Comput.* **1(1)**: 15-29.
- RAKOCEVIĆ, V. and Y. WEI. 2002. A weighted drazin inverse and applications. *Lin. Alg. and its Appl.* **350**: 25-39.
- RAKOCEVIĆ, V. and Y. WEI. 2001. The representation theory for the Drazin inverse and its applications II. *J. Austral. Math. Soc.* **70**: 189-197.
- RAO, C.R. and S.K. MITRA. 1971. *Generalized Inverses of Matrices and its Applications*. New York: Wiley.
- STEEB, W.-H. 1997. *Matrix Calculus and Kronecker Product with Applications and C++ Programs*. Singapore: World Scientific Publishing Co. Pte. Ltd.
- TIAN, Y. 1999. On the reverse order law  $(AB)_d = B_d A_d$ . *J. Math. Res. Exposition* **19(2)**: 355-358.
- WANG, G. 1997. Weighted Moore - Penrose, Drazin and Group Inverses of the Kronecker Product  $A \otimes B$ , and Some Applications. *Lin. Alg. and its Appl.* **250**: 39-50.
- WANG, G. 2002. The reverse order law for the Drazin inverses of multiple matrix products. *Lin. Alg. and its Appl.* **348**: 265-272.

- WANG, G. and Y. WEI. 1994. The iterative methods for computing the generalized inverse  $A_{MN}^+$  and  $A_{d,w}$ . *Numer. Math. J. Chinese Univ.* **16**: 366-372.
- WEI, Y. 1996. A characterization and representation of the Drazin inverse. *SIAM J. Matrix Anal. Appl.* **17**: 744-747.
- WEI, Y. and H. WU. 2000. The representation and approximation for the Drazin inverse. *J. Comput. Appl. Math.* **126**: 417-432.
- WEI, Y. 2002. A characterization for the W-Weighted Drazin inverse solution. *Appl. Math. Comput.* **125**: 303-310.
- WEI, Y. 2003. Integral representation of the W-Weighted Drazin inverse. *Appl. Math. Comput.* **144**: 3-10.
- WEI, Y., C.-W. WOO and T. LEI. 2004. A note on the perturbation of the W-Weighted Drazin inverse. *Appl. Math. Comput.* **149**: 423-430.
- WEI, Y. and D. DJORDJEVIĆ. 2003. On integral representation of the generalized inverse  $A_{T,S}^{(2)}$ . *Appl. Math. Comput.* **142**: 189-194.
- ZHANG, F. 1999. *Matrix Theory: Basic Results and Techniques*. New York: Springer-Verlag.