

Numerical Solution of Second Kind Linear Fredholm Integral Equations Using QSGS Iterative Method with High-Order Newton-Cotes Quadrature Schemes

¹Mohana Sundaram Muthuvalu and ²Jumat Sulaiman

*^{1,2}School of Science and Technology, Universiti Malaysia Sabah,
Locked Bag 2073, 88999 Kota Kinabalu, Sabah, Malaysia
E-mail: sundaram_at2@yahoo.com and jumat@ums.edu.my*

ABSTRACT

The main purpose of this paper is to examine the effectiveness of the Quarter-Sweep Gauss-Seidel (QSGS) method in solving the dense linear systems generated from the discretization of the linear Fredholm integral equations of the second kind. In addition, the applications of the various orders of closed Newton-Cotes quadrature discretization schemes will be investigated in order to form linear systems. Furthermore, the basic formulation and implementation of the proposed method are also presented. The numerical results of test examples are also included in order to verify the performance of the proposed method.

Keywords: Linear Fredholm equations, Newton-Cotes quadrature, Gauss-Seidel, Quarter-sweep iteration

INTRODUCTION

Integral equations have been one of the principal tools in various areas of science such as applied mathematics, physics, biology and engineering. On the other hand, integral equations are encountered in numerous applications in many fields including continuum mechanics, potential theory, geophysics, electricity and magnetism, kinetic theory of gases, hereditary phenomena in physics and biology, renewal theory, quantum mechanics, radiation, optimization, optimal control systems, communication theory, mathematical economics, population genetics, queuing theory, medicine, mathematical problems of radiative equilibrium, particle transport problems of astrophysics and reactor theory, acoustics, fluid mechanics, steady state heat conduction, fracture mechanics, and radiative heat transfer problems (Wang (2006)). From few types of the integral equations, the most frequently investigated integral equations are Fredholm linear equations and its nonlinear counterpart. However, in this paper, linear Fredholm integral equations of the second kind are considered.

Generally, second kind linear integral equations of Fredholm type in the generic form can be defined as follows

$$\lambda y(x) - \int_{\Gamma} K(x,t) y(t) dt = f(x), \Gamma = [a,b] \quad \lambda \neq 0 \quad (1)$$

where the parameter λ , kernel K and free term f are given, and y is the unknown function to be determined. Kernel K is called Fredholm kernel if the kernel in Equation (1) is continuous on the square $S = \{a \leq x \leq b, a \leq t \leq b\}$ or at least square integrable on this square and it is also assumed to be absolutely integrable and satisfy other properties that are sufficient to imply the Fredholm alternative theorem. Meanwhile, Equation (1) also can be rewritten in the equivalent operator form

$$(\lambda - \kappa)y = f \quad (2)$$

where the integral operator define as follows

$$\kappa y(t) = \int_{\Gamma} K(x,t) y(t) dt. \quad (3)$$

Theorem (Fredholm Alternative) (Atkinson (1997))

Let \mathcal{X} be a Banach space and let $\kappa: \mathcal{X} \rightarrow \mathcal{X}$ be compact. Then the equation $(\lambda - \kappa)y = f$, $\lambda \neq 0$ has a unique solution $x \in \mathcal{X}$ if and only if the homogeneous equation $(\lambda - \kappa)z = 0$ has only the trivial solution $z = 0$. In such a case, the operator $\lambda - \kappa: \mathcal{X} \xrightarrow[\text{onto}]{} \mathcal{X}$ has a bounded inverse $(\lambda - \kappa)^{-1}$.

Definition (Compact operators) (Atkinson (1997))

Let \mathcal{X} and Y be normed vector space and let $\kappa: \mathcal{X} \rightarrow Y$ be linear. Then κ is compact if the set $\{\kappa x \mid \|x\| \leq 1\}$ has compact closure in Y . This is equivalent to saying that for every bounded sequence $\{x_n\} \subset \mathcal{X}$, the sequences $\{\kappa x_n\}$ has a subsequence that is convergent to some points in Y . Compact operators are also called completely continuous operators.

In many application areas, numerical approaches were used widely to solve Fredholm integral equations of the second kind than the analytical method. To solve Equation (2) numerically, we either seek to determine an approximate solution by using the quadrature method (Laurie (2001); Lin (2003); Muthuvalu and Sulaiman, (2008b; 2009))

$$(\lambda I - \kappa_n) y_n = f \quad (4)$$

where I is the identity matrix and κ_n is the approximation of the κ which is obtained by discretization of κ by an n point quadrature method, or use the projection method

$$(\lambda - P_n \kappa) y_n = P_n f \quad (5)$$

where $y_n \in V_n$ and $P_n : C \rightarrow V_n$ is a projection operator in a chosen finite dimensional space V_n ; see Kaneko (1989), Chen *et al.* (2002), Maleknejad and Kajani (2003), Asady *et al.* (2005), Kajani and Vencheh (2005), Xiao *et al.* (2006), Chen *et al.* (2007), Long and Nelakanti (2007), and Oladejo *et al.* (2008). Such discretizations of integral equations lead to dense linear systems and can be prohibitively expensive to solve as n , the order of the linear system increases. Thus, iterative methods are the natural options for efficient solutions.

Consequently, the concept of the half-sweep iterative method has been proposed by Abdullah (1991) via the Explicit Decoupled Group (EDG) method to solve two-dimensional Poisson equations. Half-sweep iteration is also known as the complexity reduction approach (Hasan *et al.* (2007)). Since the implementation of half-sweep iterations will only consider half of all interior node points in a solution domain. Following to that, further studies on the applications of the half-sweep iterative methods have been reviewed by Yousif and Evans (1995), Abdullah and Ali (1996), Othman *et al.* (2000), Muthuvalu and Sulaiman (2008a; 2008b; 2009; 2011), Sulaiman *et al.* (2004a; 2007; 2008a) and Abdullah *et al.* (2006).

In 2000, Othman and Abdullah extended the concept of half-sweep iteration by introducing quarter-sweep iterative method via the Modified Explicit Group (MEG) iterative method to solve two-dimensional Poisson equations. Further studies to verify the effectiveness of the quarter-sweep iterative methods have been carried out by Othman and Abdullah (2001),

Hasan *et al.* (2005), Sulaiman *et al.* (2004b), Hasan *et al.* (2008), Sulaiman *et al.* (2008b), and Sulaiman *et al.* (2010). However, in this paper, we examined the applications of the half- and quarter-sweep iteration concepts with Gauss-Seidel (GS) iterative method by using approximation equation based on Newton-Cotes quadrature schemes for solving problem (1). The standard GS iterative method is also called as the Full-Sweep Gauss-Seidel (FSGS) method. Meanwhile, combinations of the GS method with half- and quarter-sweep iterations are called as Half-Sweep Gauss-Seidel (HSGS) and Quarter-Sweep Gauss-Seidel (QSGS) methods respectively.

The remainder of this paper is organized in following way. In next section, the formulation of the full-, half- and quarter-sweep quadrature approximation equations based on repeated Newton-Cotes schemes will be elaborated. The latter section of this paper will discuss the formulations of the FSGS, HSGS and QSGS iterative methods and some numerical results will be shown to assert the effectiveness of the proposed method. Besides that, analysis on computational complexity is also given and the concluding remarks are given in final section.


QUARTER-SWEEP QUADRATURE APPROXIMATION EQUATION

As explained in previous section, discretization method based on quadrature schemes was used to construct approximation equations for problem (1) by replacing the integral to finite sums. Generally, quadrature method can be defined as follows


$$\int_a^b y(t) dt = \sum_{j=0}^n A_j y(t_j) + \varepsilon_n(y) \quad (6)$$

where t_j ($j=0,1,2,\dots,n$) is the abscissas of the partition points of the integration interval $[a,b]$, A_j ($j=0,1,2,\dots,n$) is numerical coefficients that do not depend on the function $y(t)$ and $\varepsilon_n(y)$ is the truncation error of Equation (6). Meanwhile, Figure 1 shows the finite grid networks in order to form the full-, half- and quarter-sweep quadrature approximation equations.


Based on Figure 1, the full-, half- and quarter-sweep iterative methods will compute approximate values onto node points of type ● only until the convergence criterion is reached. According to Abdullah (1991) and, Othman and Abdullah (2000), the approximation solutions for the remaining points are calculated by using direct methods.



(a)



(b)



(c)

Figure 1: (a), (b) and (c) show distribution of uniform node points for the full-, half- and quarter-sweep cases respectively

However, in 2009, Muthuvalu and Sulaiman carried out a study to investigate the applications of the half-sweep iteration in solving dense linear system generated from the discretization of the second kind Fredholm integral equations using high-order Newton-Cotes schemes. From the results obtained, it has shown that applications of the half-sweep iteration with high-order Newton-Cotes discretization schemes reduce the accuracy of the numerical solutions and it is due to the computational technique for calculating the remaining points by using direct method.

Thus, in this paper, we will use second-order Lagrange interpolation method to compute the remaining points for both half- and quarter-sweep iterations in order to overcome the problem mentioned in Muthuvalu and Sulaiman (2009). Formulations to compute the remaining points using second order Lagrange interpolation for half- and quarter-sweep iterations are defined in Equations (7) and (8) respectively as follows

$$y_i = \begin{cases} \frac{3}{8}y_{i-1} + \frac{3}{4}y_{i+1} - \frac{1}{8}y_{i+3}, & i = 1, 3, 5, \dots, n-3 \\ \frac{3}{4}y_{i-1} + \frac{3}{8}y_{i+1} - \frac{1}{8}y_{i-3}, & i = n-1 \end{cases} \quad (7)$$

$$y_i = \begin{cases} \frac{3}{8}y_{i-2} + \frac{3}{4}y_{i+2} - \frac{1}{8}y_{i+6}, & i = 2, 6, 10, \dots, n-6 \\ \frac{3}{4}y_{i-2} + \frac{3}{8}y_{i+2} - \frac{1}{8}y_{i-6}, & i = n-2 \\ \frac{3}{8}y_{i-1} + \frac{3}{4}y_{i+1} - \frac{1}{8}y_{i+3}, & i = 1, 3, 5, \dots, n-3 \\ \frac{3}{4}y_{i-1} + \frac{3}{8}y_{i+1} - \frac{1}{8}y_{i-3}, & i = n-1 \end{cases} \quad (8)$$

By applying Equation (6) into Equation (1) and neglecting the error, $\epsilon_n(y)$, a system of linear equations can be formed for approximation values of $y(t)$. The following linear system generated using quadrature method can be easily shown in matrix form as follows

$$M y = f \quad (9)$$

where

$$M = \begin{bmatrix} \lambda - A_0 K_{0,0} & -A_p K_{0,p} & -A_{2p} K_{0,2p} & \dots & -A_n K_{0,n} \\ -A_0 K_{p,0} & \lambda - A_p K_{p,p} & -A_{2p} K_{p,2p} & \dots & -A_n K_{p,n} \\ -A_0 K_{2p,0} & -A_p K_{2p,p} & \lambda - A_{2p} K_{2p,2p} & \dots & -A_n K_{2p,n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -A_0 K_{n,0} & -A_p K_{n,p} & -A_{2p} K_{n,2p} & \dots & \lambda - A_n K_{n,n} \end{bmatrix} \left(\left(\binom{n}{p} + 1 \right) \times \left(\binom{n}{p} + 1 \right) \right),$$

$$\begin{aligned} \underline{y} &= [y_0 \quad y_p \quad y_{2p} \quad \cdots \quad y_{n-2p} \quad y_{n-p} \quad y_n]^T, \\ \underline{f} &= [f_0 \quad f_p \quad f_{2p} \quad \cdots \quad f_{n-2p} \quad f_{n-p} \quad f_n]^T. \end{aligned}$$

In order to facilitate the formulation of the full-, half- and quarter-sweep quadrature approximation equations for problem (1), further discussion will be restricted onto Newton-Cotes quadrature method, which is based on interpolation formulas with equally spaced data. In this paper, three different schemes in Newton-Cotes quadrature method such as repeated trapezoidal (RT), repeated Simpson's $\frac{1}{3}$ (RS1) and repeated Simpson's $\frac{3}{8}$ (RS2) schemes will be applied to discretize the problem (1). RT, RS1 and RS2 are first, second and third order schemes respectively. Further discussions on Newton-Cotes quadrature method to solve Fredholm integral equations can be found in Atkinson (1997), and Muthuvalu and Sulaiman (2009).

Based on RT, RS1 and RS2 schemes, numerical coefficients A_j will satisfy following relations respectively.

$$A_j = \begin{cases} \frac{1}{2}ph, & j = 0, n \\ ph, & otherwise \end{cases} \quad (10)$$

$$A_j = \begin{cases} \frac{1}{3}ph, & j = 0, n \\ \frac{4}{3}ph, & j = p, 3p, 5p, \dots, n-p \\ \frac{2}{3}ph, & otherwise \end{cases} \quad (11)$$

$$A_j = \begin{cases} \frac{3}{8}ph, & j = 0, n \\ \frac{3}{4}ph, & j = 3p, 6p, 9p, \dots, n - 3p \\ \frac{9}{8}ph, & \text{otherwise} \end{cases} \quad (12)$$

where the constant step-size, h is defined as follows

$$h = \frac{b-a}{n} \quad (13)$$

and n is the number of subintervals in the interval $[a, b]$. Meanwhile, the value of p , which corresponds to 1, 2 and 4, represents the full-, half- and quarter-sweep cases respectively.

FORMULATION OF THE ITERATIVE METHODS

As afore-mentioned, FSGS, HSGS and QSGS iterative methods will be applied to solve linear system generated from the discretization of the problem (1), as shown in Equation (9). Let matrix M be decomposed into

$$M = D - L - U \quad (14)$$

where D , $-L$ and $-U$ are diagonal, strictly lower triangular and strictly upper triangular matrices respectively. Thus, the general scheme for FSGS, HSGS and QSGS iterative methods can be written as

$$\underline{y}^{(k+1)} = (D - L)^{-1} \left(U \underline{y}^{(k)} + \underline{f} \right). \quad (15)$$

Actually, the iterative methods attempt to find a solution by repeatedly solving the linear system using approximations to the vector \underline{y} and continue until the solution is within a predetermined acceptable bound on the error. Based on Abdullah (1991) and, Othman and Abdullah (2000), the general algorithm for FSGS, HSGS and QSGS iterative methods to solve problem (1) would be generally described in Algorithm 1.

Algorithm 1: FSGS, HSGS and QSGS methods

For $i = 0, p, 2p, \dots, n-2p, n-p, n$ and $j = 0, p, 2p, \dots, n-2p, n-p, n$ calculate

$$y_i^{(k+1)} \leftarrow \begin{cases} \left(f_i + \sum_{j=p}^n A_j K_{i,j} y_j^{(k)} \right) / \lambda - A_i K_{i,i} , & i = 0 \\ \left(f_i + \sum_{j=0}^{n-p} A_j K_{i,j} y_j^{(k+1)} \right) / \lambda - A_i K_{i,i} , & i = n \\ \left(f_i + \sum_{j=0}^{i-p} A_j K_{i,j} y_j^{(k+1)} + \sum_{j=i+p}^n A_j K_{i,j} y_j^{(k)} \right) / \lambda - A_i K_{i,i} , & i = \text{otherwise} \end{cases}$$

NUMERICAL SIMULATIONS

In order to compare the performances of the iterative methods, several experiments were carried out on the following Fredholm integral equations problems.

Example 1 (Wang (2006))

Consider the integral equation

$$y(x) - \int_0^1 (4xt - x^2) y(t) dt = x \tag{16}$$

and the exact solution of problem (16) is given by $y(x) = 24x - 9x^2$.

Example 2 (Polyanin and Manzhirov (1998))

Consider the integral equation

$$y(x) - \int_0^1 (x^2 + t^2) y(t) dt = x^6 - 5x^3 + x + 10. \tag{17}$$

Exact solution of problem (17) is

$$y(x) = x^6 - 5x^3 + \frac{1045}{28}x^2 + x + \frac{2141}{84}.$$

There are three parameters considered in numerical comparison such as number of iterations, execution time and maximum absolute error. Throughout the experiments, the convergence test considered the tolerance error of the $\epsilon = 10^{-10}$. The experiments were carried out on several different mesh sizes such as 240, 480, 960, 1920, 3840 and 7680. Results of numerical simulations, which were obtained from implementations of the FSGS, HSGS and QSGS iterative methods for Examples 1 and 2, have been recorded in Tables 1 and 2 respectively.

TABLE 1: Comparison of a number of iterations, execution time (seconds) and maximum absolute error for the iterative methods for Example 1

Mesh Size	Methods	Number of iterations			Execution time (seconds)			Maximum absolute error		
		FSGS	HSGS	QSGS	FSGS	HSGS	QSGS	FSGS	HSGS	QSGS
240	RT	193	192	189	0.51	0.20	0.19	2.136 E-3	8.547 E-3	3.425 E-2
	RS1	193	191	188	0.52	0.21	0.21	7.262 E-10	7.488 E-10	7.055 E-10
	RS2	193	192	189	0.54	0.26	0.23	7.376 E-10	6.830 E-10	6.641 E-10
480	RT	194	193	192	1.82	0.63	0.24	5.339 E-4	2.136 E-3	8.547 E-3
	RS1	194	193	191	1.84	0.66	0.29	7.156 E-10	7.262 E-10	7.488 E-10
	RS2	194	193	192	1.86	0.73	0.32	7.212 E-10	7.376 E-10	6.830 E-10
960	RT	194	194	193	6.72	2.01	0.85	1.335 E-4	5.339 E-4	2.136 E-3
	RS1	194	194	193	6.79	2.02	0.94	7.552 E-10	7.156 E-10	7.262 E-10
	RS2	194	194	193	6.84	2.25	0.97	7.581 E-10	7.212 E-10	7.376 E-10
1920	RT	195	194	194	26.24	7.58	2.63	3.337 E-5	1.335 E-4	5.339 E-4
	RS1	195	194	194	26.93	7.61	2.75	6.868 E-10	7.552 E-10	7.156 E-10
	RS2	195	194	194	27.05	7.79	2.77	8.341 E-6	7.581 E-10	7.212 E-10
3840	RT	195	195	194	100.09	27.48	8.55	6.961 E-10	3.337 E-5	1.335 E-4
	RS1	195	195	194	101.99	28.08	8.98	6.868 E-10	6.868 E-10	7.552 E-10
	RS2	195	195	194	102.30	29.33	9.09	6.968 E-10	6.881 E-10	7.581 E-10
7680	RT	195	195	195	397.81	108.77	31.44	2.085 E-6	8.341 E-6	3.337 E-5
	RS1	195	195	195	402.91	110.17	33.37	7.008 E-10	6.961 E-10	6.868 E-10
	RS2	195	195	195	403.95	111.43	34.09	7.012 E-10	6.968 E-10	6.881 E-10

TABLE 2: Comparison of a number of iterations, execution time (seconds) and maximum absolute error for the iterative methods for Example 2

Mesh Size	Methods	Number of iterations			Execution time (seconds)			Maximum absolute error		
		FSGS	HSGS	QSGS	FSGS	HSGS	QSGS	FSGS	HSGS	QSGS
240	RT	56	55	55	0.18	0.14	0.09	2.174 E-3	8.697 E-3	3.481 E-2
	RS1	56	55	55	0.20	0.16	0.10	1.124 E-8	3.342 E-7	2.742 E-6
	RS2	56	55	55	0.21	0.17	0.11	2.542 E-8	3.567 E-7	3.102 E-6
480	RT	56	56	55	0.56	0.23	0.16	5.435 E-4	2.174 E-3	8.697 E-3
	RS1	56	56	55	0.57	0.27	0.17	5.882 E-10	4.122 E-8	3.342 E-7
	RS2	56	56	55	0.59	0.29	0.19	1.474 E-9	4.264 E-8	3.567 E-7
960	RT	56	56	56	2.52	0.58	0.34	1.359 E-4	5.435 E-4	2.174 E-3
	RS1	56	56	56	2.57	0.63	0.38	8.305 E-11	5.108 E-10	4.122 E-8
	RS2	56	56	56	2.65	0.64	0.40	3.887 E-11	5.196 E-9	4.264 E-8
1920	RT	56	56	56	8.24	2.71	1.11	3.397 E-5	1.359 E-4	5.435 E-4
	RS1	56	56	56	8.40	2.79	1.17	1.260 E-10	6.410 E-11	5.108 E-10
	RS2	56	56	56	8.71	2.83	1.20	1.227 E-10	6.355 E-11	5.196 E-9
3840	RT	56	56	56	31.62	11.04	5.40	8.492 E-6	3.397 E-5	1.359 E-4
	RS1	56	56	56	32.04	11.87	5.77	1.301 E-10	9.188 E-10	6.410 E-11
	RS2	56	56	56	34.01	12.04	5.91	1.299 E-10	9.155 E-10	6.355 E-11
7680	RT	56	56	56	122.70	35.16	22.83	2.123 E-6	8.492 E-6	3.397 E-5
	RS1	56	56	56	124.66	36.79	24.89	1.309 E-10	2.291 E-10	9.188 E-10
	RS2	56	56	56	127.33	38.14	25.11	1.310 E-10	2.291 E-10	9.155 E-10

COMPUTATIONAL COMPLEXITY ANALYSIS

In order to measure the computational complexity of the iterative methods, an estimation of the amount of the computational works required for both methods have been conducted. The computational works are estimated by considering the arithmetic operations performed per iteration.

Based on Algorithm 1, it can be observed that there are $\binom{\frac{n}{p}+1}{p}$

additions/subtractions (ADD/SUB) and $2\binom{\frac{n}{p}+1}{p}$ multiplications/divisions

(MUL/DIV) in computing a value for each node point in the solution domain. From the order of the coefficient matrix, M in Equation (9), the

total number of arithmetic operations per iteration for the FSGS, HSGS and QSGS iterative methods has been summarized in Table 3.

TABLE 3: Total number of arithmetic operations per iteration for FSGS, HSGS and QSGS methods

Methods	Arithmetic Operation	
	ADD/SUB	MUL/DIV
FSGS	$(n+1)^2$	$2(n+1)^2$
HSGS	$\left(\frac{n}{2}+1\right)^2$	$2\left(\frac{n}{2}+1\right)^2$
QSGS	$\left(\frac{n}{4}+1\right)^2$	$2\left(\frac{n}{4}+1\right)^2$

CONCLUSIONS

In this paper, we present applications of the half- and quarter-sweep iterative methods for solving dense linear systems arising from the discretization of the second kind linear Fredholm integral equations by using three different orders of Newton-Cotes quadrature discretization schemes such as RT, RS1 and RS2 schemes. It has shown that the quadrature approximation equations based on Newton-Cotes schemes can be easily formulated and rewritten in general form as shown in Equation (9).

Through numerical results obtained for both Examples 1 and 2 (refer Tables 1 and 2), it shows that number of iterations for HSGS and QSGS methods are nearly same compared to the FSGS method. Through the observation in Tables 1 and 2, HSGS and QSGS iterative methods reduce the execution time compared to the FSGS method. Computational time for FSGS, HSGS and QSGS iterative methods with RS1 and RS2 schemes are increased compared to the iterative methods with RT scheme. It is due to the computational complexity of the high-order discretization schemes. In terms of accuracy of numerical solutions obtained, RS1 and RS2 schemes are more accurate than the RT scheme. Besides that, applications of second order Lagrange interpolation to compute remaining points managed to overcome the problem discussed in Muthuvalu and Sulaiman (2009).

Overall, the numerical results show that the QSGS method is a better method compared to the FSGS and HSGS methods in the sense of number of iterations and execution time. This is mainly because of computational

complexity of the QSGS method which is approximately 50% and 75% less than HSGS and FSGS methods respectively (refer Table 3).

ACKNOWLEDGEMENT

The authors acknowledge the Postgraduate Research Grant, Universiti Malaysia Sabah (GPS0003-SG-1/2009) for the completion of this article.

REFERENCES

- Abdullah, A. R. 1991. The four point Explicit Decoupled Group (EDG) method: A fast Poisson solver. *Int. J. Comput. Math.*, **38**: 61-70.
- Abdullah, A. R. and Ali, N. H. M. 1996. A comparative study of parallel strategies for the solution of elliptic pde's. *Parallel Algorithms and Applications*, **10**: 93-103.
- Abdullah, M. H., Sulaiman, J. and Othman, A. 2006. A numerical assessment on water quality model using the Half-Sweep Explicit Group methods. *Gading*, **10**: 99-110.
- Asady, B., Kajani, M. T., Vencheh, A. H. and Heydari, A. 2005. Solving second kind integral equations with hybrid Fourier and block-pulse functions. *Appl. Math. Comput.*, **160**: 517-522.
- Atkinson, K. E. 1997. *The Numerical Solution of Integral Equations of the Second Kind*, United Kingdom: Cambridge University Press.
- Chen, Z., Micchelli, C. A. and Xu, Y. 2002. Fast collocation methods for second kind integral equations. *SIAM J. Numer. Anal.*, **40**(1): 344-375.
- Chen, Z., Wu, B. and Xu, Y. 2007. Fast numerical collocation solutions of integral equations. *Commun. Pure Appl. Anal.*, **6**(3): 643-666.
- Hasan, M.K., Othman, M., Abbas, Z., Sulaiman, J. and Ahmad, F. 2007. *Parallel solution of high speed low order FDTD on 2D free space wave propagation*. In O. Gervasi and M. Gavrilova (Eds.),

Computational Science and Its Applications, Lecture Notes in Computer Science (LNCS 4706): 13-24. Springer-Verlag, Berlin.

- Hasan, M. K., Othman, M., Johari, R., Abbas, Z. and Sulaiman, J. 2005. The HSLO(3)-FDTD with direct-domain and temporary-domain approaches on infinite space wave propagation. In B.M. Ali, M. Ismail & H. Mohamad, *Proceedings of the 13th IEEE International Conference on Network*: 1002-1007.
- Hasan, M. K., Sulaiman, J. and Othman, M. 2008. Implementation of red black strategy to quarter-sweep iteration for solving first order hyperbolic equations. In H.B. Zaman, T.M.T. Sembok, K.v. Rijsbergen, L. Zadeh, P. Bruza, T. Shih and M. N. Taib, *Proceedings of the International Symposium on Information Technology* : 1864-1869.
- Kajani, M. T. and Vencheh, A. H. 2005. Solving second kind integral equations with Hybrid Chebyshev and Block-Pulse functions. *Appl. Math. Comput.*, **163**: 71-77.
- Kaneko, H. 1989. A projection method for solving Fredholm integral equations of the second kind. *Appl. Numer. Math.*, **5**(4): 333-344.
- Laurie, D. P. 2001. Computation of Gauss-type quadrature formulas. *J. Comput. Appl. Math.*, **127**: 201-217.
- Lin, F. R. 2003. Preconditioned iterative methods for the numerical solution of Fredholm equations of the second kind. *Calcolo*, **40**: 231-248.
- Long, G. and Nelakanti, G. 2007. Iteration methods for Fredholm integral equations of the second kind. *Comput. Math. Appl.*, **53**: 886-894.
- Maleknejad, K. and Kajani, M.T. 2003. Solving second kind integral equations by Galerkin methods with hybrid Legendre and Block-Pulse functions. *Appl. Math. Comput.*, **145**: 623-629.
- Muthuvalu, M. S. and Sulaiman, J. 2008a. Half-Sweep Geometric Mean method for solution of linear Fredholm equations. *Matematika*, **24**(1): 75-84.

- Muthuvalu, M. S. and Sulaiman, J. 2008b. Numerical solutions of second kind Fredholm integral equations using Half-Sweep Geometric Mean method. In H.B. Zaman, T.M.T. Sembok, K.v. Rijsbergen, L. Zadeh, P. Bruza, T. Shih and M. N. Taib, *Proceedings of the International Symposium on Information Technology: 1927-1934*.
- Muthuvalu, M. S. and Sulaiman, J. 2009. Half-Sweep Arithmetic Mean method with high-order Newton-Cotes quadrature schemes to solve linear second kind Fredholm equations. *Journal of Fundamental Sciences*, **5**(1): 7-16.
- Muthuvalu, M. S. and Sulaiman, J. 2011. Half-Sweep Arithmetic Mean method with composite trapezoidal scheme for solving linear Fredholm integral equations. *Appl. Math. Comput.*, In Press. DOI: 10.1016/j.amc.2010.12.013
- Oladejo, S. O., Mojeed, T.A. and Olurode, K.A. 2008. The application of cubic spline collocation to the solution of integral equations. *J. Appl. Sci. Res.*, **4**(6): 748-753.
- Othman, M. and Abdullah, A.R. 2000. An efficient Four Points Modified Explicit Group Poisson solver. *Int. J. Comput. Math.*, **76**: 203-217.
- Othman, M. and Abdullah, A. R. 2001. *Implementation of the Parallel Four Points Modified Explicit Group Iterative Algorithm on Shared Memory Parallel Computer*. In V. Malyskin (Ed.), *Parallel Computing Technologies, Lecture Notes in Computer Science (LNCS 2127)*: 480-489. Springer-Verlag, Berlin.
- Othman, M., Sulaiman, J. and Abdullah, A.R. 2000. A parallel halfsweep multigrid algorithm on the shared memory multiprocessors. *Malaysian Journal of Computer Science*, **13**(2): 1-6.
- Polyanin, A.D. and Manzhirov, A.V. 1998. *Handbook of Integral Equations*, Florida: CRC Press LLC.
- Sulaiman, J., Hasan, M. K. and Othman, M. 2004a. *The Half-Sweep Iterative Alternating Decomposition Explicit (HSIADE) method for diffusion equation*. In J. Zhang, J.-H. He and Y. Fu (Eds.), *Computational and Information Science, Lecture Notes in Computer Science (LNCS 3314)*: 57-63. Springer-Verlag, Berlin.

- Sulaiman, J., Hasan, M. K. and Othman, M. 2007. *Red-Black Half-Sweep iterative method using triangle finite element approximation for 2D Poisson equations*. In Y. Shi *et al.* (Eds.), Computational Science, Lecture Notes in Computer Science (LNCS 4487): 326-333. Springer-Verlag, Berlin.
- Sulaiman, J., Hasan, M. K. and Othman, M. 2010. MEGSOR iterative method for the triangle element solution of 2D Poisson equations. *Procedia Computer Science*, **1**: 377-385.
- Sulaiman, J., Othman, M. and Hasan, M. K. 2004b. Quarter-Sweep Iterative Alternating Decomposition Explicit algorithm applied to diffusion equations. *Int. J. Comput. Math.*, **81**(12): 1559-1565.
- Sulaiman, J., Othman, M. and Hasan, M. K. 2008a. *Half-Sweep Algebraic Multigrid (HSAMG) method applied to diffusion equations*. In Modeling, Simulation and Optimization of Complex Processes: 547-556. Springer-Verlag, Berlin.
- Sulaiman, J., Saudi, A., Abdullah, M. H., Hasan, M. K. and Othman, M. 2008b. Quarter-Sweep Arithmetic Mean algorithm for water quality model. In H.B. Zaman, T.M.T. Sembok, K.V. Rijsbergen, L. Zadeh, P. Bruza, T. Shih and M. N. Taib, *Proceedings of the International Symposium on Information Technology*: 1859-1863.
- Wang, W. 2006. A new mechanical algorithm for solving the second kind of Fredholm integral equation. *Appl. Math. Comput.*, **172**: 946-962.
- Xiao, J. Y., Wen, L. H. and Zhang, D. 2006. Solving second kind Fredholm integral equations by periodic wavelet Galerkin method. *Appl. Math. Comput.*, **175**: 508-518.
- Yousif, W. S. and Evans, D. J. 1995. Explicit De-coupled Group iterative methods and their implementations. *Parallel Algorithms and Applications*, **7**: 53-71.