



The Interval Zero-Symmetric Single-Step IZSS1-5D for the Simultaneous Bounding of Real Polynomial Zeros

¹Atiyah Wan Mohd Sham, ¹Mansor Monsi and ^{2*}Nasruddin Hassan

¹*Department of Mathematics, Faculty of Science,
Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia*

²*School of Mathematical Sciences, Faculty of Science and Technology,
Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia*

E-mail: nas@ukm.edu.my

*Corresponding author

ABSTRACT

A new modified method IZSS1-5D for the simultaneously bounding all the real zeros of a polynomial is formulated in this paper. The efficiency of this method is measured on the CPU times and the number of iterations after satisfying the convergence criteria where the results are obtained using five tested polynomials. The analysis performed shows that the R -order of convergence of this new procedure is at least five. The programming language used to obtain the numerical results is Matlab R2012, a software in association with Intlab V5.5 toolbox. The numerical results indicate that the procedure IZSS1-5D outperformed the IZSS1 in computational times and number of iterations.

Keywords: CPU time, initial disjoint intervals, interval, number of iteration.

1. INTRODUCTION

Consider $p : R^1 \rightarrow R^1$ a polynomial of degree $n > 1$ defined by

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_0 = \prod_{j=1}^n (x - x_j^*) \quad (1)$$

where $a_i \in R^1 (i=1, \dots, n)$ are given. Suppose that p has n distinct values $x_i^* \in R (i=1, \dots, n)$ and that $X_i^{(0)} \in I(R)$ (set of real intervals) $(i=1, \dots, n)$ are such that

$$x_i^* \in X_i^{(0)}, (i=1, \dots, n) \tag{2}$$

and

$$X_i^{(0)} \cap X_j^{(0)} = \emptyset, (i, j=1, \dots, n; i \neq j). \tag{3}$$

There are researchers who came out with the new idea of applying the interval arithmetic, such as Moore, 1979 and Alefeld and Herzberger, 1983 in determining the convergence rate of the interval procedures related to bounding polynomial zeros. These were then followed by Bakar et al., 2012, Sham et al., 2013a; 2013b, Jamaludin et al., 2013a; 2013b; 2013c; 2013d; 2014a; 2014b and Rusli et al., 2014a; 2014b. Interval iterative procedure for simultaneous inclusion of simple polynomial zeros determine bounded closed intervals which contain an exact polynomial zeros. It can be used to determine very narrow computationally rigorous bound on polynomial zeros. It is a very significant way of obtaining reliable bounds on the zeros as the intervals sequences generated by the procedures always converges to the zeros.

The aim of this paper is to present the interval zero symmetric single-step procedure, IZSS1-5D which is the modification of interval zero symmetric single-step procedure, IZSS1 (Rusli et al., 2011). IZSS1-5D method has been modified in order to increase the efficiency of the method. We repeat the same inner looping and add up another step to test the accuracy. The efficiency of the algorithm is measured numerically by taking the CPU time and also the number of iterations of the algorithm.

2. THE INTERVAL ZORO SYMMETRIC SINGLE-STEP PROCEDURE IZSS1-5D

An extension of the interval single-step procedure IS is the interval symmetric single-step procedure ISS. This idea is basically ideas of Alefeld and Herzberger, 1983 and Monsi and Wolfe, 1988. The procedure IZSS1-5D consists of generating the interval sequence $\{X_i^{(k)}\}, (i=1, \dots, n)$ from the following algorithm,

Step 1: Set $k = 0$, (4a)

Step 2: For $k \geq 0, x_i^{(k)} = \text{mid}(X_i^{(k)}), (i=1, \dots, n);$ (4b)

$$\text{Step 3: Let } \delta_j = \delta_{ij}^{(k)} = -p(x_i^{(k)}) / \prod_{j \neq i}^n (x_i^{(k)} - x_j^{(k)}) \quad (4c)$$

Step 4:

$$X_i^{(k,1)} = \left\{ x_i^{(k)} - \frac{p(x_i^{(k)})}{\left\{ \prod_{j=1}^{i-1} (x_i^{(k)} - X_j^{(k,1)}) \prod_{j=i+1}^n (x_i^{(k)} - X_j^{(k,0)} - 5\delta_j) \right\}} \right\} \cap X_i^{(k)} \quad (4d)$$

(i = 1, ..., n)

Step 5:

$$X_i^{(k,2)} = \left\{ x_i^{(k)} - \frac{p(x_i^{(k)})}{\left\{ \prod_{j=1}^{i-1} (x_i^{(k)} - X_j^{(k,1)}) \prod_{j=i+1}^n (x_i^{(k)} - X_j^{(k,2)}) \right\}} \right\} \cap X_i^{(k,1)} \quad (4e)$$

(i = n, ..., 1)

Step 6:

$$X_i^{(k,3)} = \left\{ x_i^{(k)} - \frac{p(x_i^{(k)})}{\left\{ \prod_{j=1}^{i-1} (x_i^{(k)} - X_j^{(k,3)}) \prod_{j=i+1}^n (x_i^{(k)} - X_j^{(k,2)}) \right\}} \right\} \cap X_i^{(k,2)} \quad (4f)$$

(i = 1, ..., n)

$$\text{Step 7: } X_i^{(k+1)} = X_i^{(k+1,0)} = X_i^{(k,3)} \quad (4g)$$

Step 8: If the widths of intervals $X_i^{(k+1)} < \varepsilon$ (i = 1, ..., n) then stop, else set $k = k + 1$ and go to Step 2.

The procedure IZSS1-5D has the following attractive features:

- The use of $5\delta_j$ ($j = 1, \dots, n$) in (4d) will improve the efficiency of this procedure.
- The values $p(x_i^{(k)})(i = 1, \dots, n)$ which are computed for use in (4d) are re-used in (4e) and (4f).
- The products $\prod_{j=1}^{i-1} (x_i^{(k)} - x_j^{(k,1)})$ ($i = 2, \dots, n$) which are computed for use in (4d) are re-used in (4e).
- The products $\prod_{j=1}^{i-1} (x_i^{(k)} - x_j^{(k,2)})$ ($i = 2, \dots, n$) which are computed for use in (4e) are re-used in (4f).

The above procedure without $5\delta_j$ ($j = 1, \dots, n$) in (4d) is the procedure IZSS1 in Rusli et al., 2011. It is an extension of the procedure ISS1-5D (Sham et al., 2013a).

3. THE RATE OF CONVERGENCE OF IZSS1-5D

The following theorem contains the conditions so that the algorithm in (4a)-(4g) converges to the zeros of the polynomial in (1).

Theorem 3.1: If (i) (2) and (3) hold; (ii) the sequences $\{X_i^{(k)}\}(i = 1, \dots, n)$ are generated from (4), then $(\forall k \geq 0) x_i^* \in X_i^{(k+1)} \subseteq X_i^{(k)} (i = 1, \dots, n)$. If also (iii) $0 \notin D_i \in I(R)$ is such that $p'(x) \in D_i (\forall x \in X_i^{(0)}) (i = 1, \dots, n)$, then $X_i^{(k)} \rightarrow x_i^* (k \rightarrow \infty)$ and $(\forall k \geq 0) (i = 1, \dots, n)$

$$w(X_i^{(k+1)}) \leq \frac{1}{2} \left(1 - \frac{d_H}{d_{IS}} \right) w(X_i^{(k)}), \tag{5}$$

where $w(X_i^{(k)}) = w([x_{iH}^{(k)}, x_{iS}^{(k)}]) = x_{iS}^{(k)} - x_{iH}^{(k)}$. Hence, $O_R(IZSS1-5D, x_i^*) \geq 5$.

Proof.

The proof that $x_i^* \in X_i^{(k+1)} \subseteq X_i^{(k)} (i = 1, \dots, n) (\forall k \geq 0)$ holds is almost identical with the corresponding proofs in Monsi and Wolfe, 1988 and Ortega and Rheinboldt, 1970 and is therefore omitted.

As the proof in Monsi et al., 2012, it may be shown that $\exists \alpha > 0$ such that $(\forall k \geq 0)$,

$$w_i^{(k,1)} \leq \beta w_i^{(k,0)} \left\{ \sum_{j=1}^{i-1} w_j^{(k,1)} + \sum_{j=i+1}^n (w_j^{(k,0)})^2 \right\} \quad (i = 1, \dots, n), \quad (6)$$

$$w_i^{(k,2)} \leq \beta w_i^{(k,0)} \left\{ \sum_{j=1}^{i-1} w_j^{(k,1)} + \sum_{j=i+1}^n w_j^{(k,2)} \right\} \quad (i = n, \dots, 1), \quad (7)$$

and

$$w_i^{(k,3)} \leq \beta w_i^{(k,0)} \left\{ \sum_{j=1}^{i-1} w_j^{(k,3)} + \sum_{j=i+1}^n w_j^{(k,2)} \right\} \quad (i = 1, \dots, n), \quad (8)$$

where

$$w_i^{(k,s)} = (n-1) \alpha w (X_i^{(k,s)}) \quad (s = 0, 1, 2), \quad (9)$$

and

$$\beta = \frac{1}{n-1} . \quad (10)$$

Let

$$u_i^{(1,1)} = \begin{cases} 3 & (i = 1, \dots, n-1) \\ 4 & (i = n) \end{cases} \quad (11)$$

$$u_i^{(1,2)} = \begin{cases} 5 & (i = 1) \\ 4 & (i = 2, \dots, n) \end{cases} \quad (12)$$

and

$$u_i^{(1,3)} = \begin{cases} 5 & (i = 1, \dots, n-1) \\ 6 & (i = n) \end{cases} \quad (13)$$

and for $(r = 1, 2, 3)$, let

$$u_i^{(k+1,r)} = \begin{cases} 5u_i^{(k,r)} & (i = 1, \dots, n-1) \\ 5u_i^{(k,r)} + 1 & (i = n) \end{cases} , \quad (14)$$

Then by (11) - (14), for $(\forall k \geq 0)$

$$u_i^{(k,1)} = \begin{cases} 3(5^{(k-1)}) & (i = 1, \dots, n-1) \\ \frac{17}{4}(5^{(k-1)}) - \frac{1}{4} & (i = n) \end{cases} , \quad (15)$$

$$u_i^{(k,2)} = \begin{cases} 5(5^{(k-1)}) & (i = 1) \\ 4(5^{(k-1)}) & (i = 2, \dots, n-1) , \\ \frac{17}{4}(5^{(k-1)}) - \frac{1}{4} & (i = n) \end{cases} \quad (16)$$

and

$$u_i^{(k,3)} = \begin{cases} 5(5^{(k-1)}) & (i = 1, \dots, n-1) \\ \frac{25}{4}(5^{(k-1)}) - \frac{1}{4} & (i = n) \end{cases} , \quad (17)$$

Suppose, without any loss of generality, that

$$w_i^{(0,0)} \leq h < 1 \quad (i = 1, \dots, n) \quad (18)$$

Then by inductive argument it follows from (9) - (18) that for $(i = 1, \dots, n)$
 $(k \geq 0)$

$$w_i^{(k,1)} \leq h^{u_i^{(k+1,1)}}, \quad w_i^{(k,2)} \leq h^{u_i^{(k+1,2)}} \quad \text{and} \quad w_i^{(k,3)} \leq h^{u_i^{(k+1,3)}}, \quad (19)$$

whence by (17)-(19) and (4g),

$$w_i^{(k+1)} \leq h^{5^{(k+1)}} \quad (i = 1, \dots, n) . \quad (20)$$

So, $(\forall k \geq 0)$, by (9) and (20),

$$w(X_i^{(k)}) \leq \left(\frac{\beta}{\alpha}\right)h^{5^{(k)}} \quad (i = 1, \dots, n) , \quad \alpha > 0. \quad (21)$$

Let

$$w^{(k)} = \max_{1 \leq i \leq n} \{w(X_i^{(k)})\} \quad (22)$$

Then by (21) and (22),

$$w^{(k)} \leq \left(\frac{\beta}{\alpha}\right)h^{5^k} \quad (\forall k \geq 0). \quad (23)$$

So, by (23), and the definition of R -factor in Ortega and Rheinboldt, 1970, we have

$$\begin{aligned} R_5(w^{(k)}) &= \limsup_{k \rightarrow \infty} \{ (w^{(k)})^{1/(5^k)} \} \\ &= \lim_{k \rightarrow \infty} \left\{ \left(\frac{\beta}{\alpha} \right)^{1/(5^k)} h \right\} \\ &= h < 1. \end{aligned}$$

Therefore, it is proven that according to Alefeld and Herzberger, 1983, Monsi and Wolfe, 1988 and Ortega and Rheinboldt, 1970, the order of convergence of IZSS1-5D is at least five or $O_R(\text{IZSS1-5D}, x_i^*) \geq 5$, ($i = 1, \dots, n$).

4. RESULTS AND DISCUSSIONS

We use the Intlab V5.5 toolbox of Rump, 1999 for Matlab R2012 to get the following results below. The algorithms IZSS1 and IZSS1-5D are run on five test polynomials where the stopping criterion used is $w^{(k)} \leq 10^{-10}$. The polynomials are as follows.

Test polynomial 1 : (Sham et al., 2013c)

$$p = \lambda^6 - 44\lambda^4 + 453\lambda^2 - 990$$

where

$$\begin{aligned} n &= 6, \quad a_1 = 5.4772, \quad a_2 = -5.4772, \quad a_3 = 3.3166, \quad a_4 = 1.7321, \\ a_5 &= -3.3166, \quad a_6 = -1.7321. \\ b_i &= 1 \quad (i = 1, \dots, 5) \end{aligned}$$

Initial intervals:

$$\begin{aligned} X_1^{(0)} &= [1, 2], \quad X_2^{(0)} = [3, 4], \quad X_3^{(0)} = [5, 6], \quad X_4^{(0)} = [-2, -1], \\ X_5^{(0)} &= [-4, -3], \quad X_6^{(0)} = [-6, -5]. \end{aligned}$$

Test polynomial 2 : (Ehrlic, 1967, Alefeld and Herzberger, 1983)

$$p = \lambda^5 - 35.6\lambda^4 + 482.86\lambda^3 - 3090.376\lambda^2 + 9197.7665\lambda - 9931.285$$

where

$$n = 5, a_1 = 11.5, a_2 = 9.1, a_3 = 7.3, a_4 = 5.2, a_5 = 2.5, b_i = 1 (i = 1, \dots, 4)$$

Initial intervals:

$$X_1^{(0)} = [-2.5, 2.1], X_2^{(0)} = [2.2, 4.5], X_3^{(0)} = [4.6, 7.9], \\ X_4^{(0)} = [8.0, 10.8], X_5^{(0)} = [10.9, 13.1]$$

Test polynomial 3: (Sham et al., 2013c)

$$p = \lambda^4 - 10\lambda^3 + 35\lambda^2 - 50\lambda + 24$$

where

$$n = 4, a_1 = 1, a_2 = 2, a_3 = 3, a_4 = 4, b_i = 1 (i = 1, \dots, 3)$$

Initial intervals:

$$X_1^{(0)} = [0.6, 1.3], X_2^{(0)} = [1.6, 2.3], X_3^{(0)} = [2.6, 3.3], X_4^{(0)} = [3.6, 4.3].$$

Test polynomial 4: (Alefeld and Herzberger, 1983)

$$p = \lambda^9 - 398\lambda^7 + 45944\lambda^5 - 1778055\lambda^3 + 17863791\lambda$$

where

$$n = 9, a_1 = 15, a_2 = 10, a_3 = 7, a_4 = 4, a_5 = 0, a_6 = -4, a_7 = -7, \\ a_8 = -10, a_9 = -15, b_i = 1 (i = 1, \dots, 8)$$

Initial intervals:

$$X_1^{(0)} = [12.0, 17.0], X_2^{(0)} = [8.6, 11.2], X_3^{(0)} = [5.2, 8.4], X_4^{(0)} = [2.4, 5.0], \\ X_5^{(0)} = [-2.0, 2.2], X_6^{(0)} = [-6.4, -2.9], \\ X_7^{(0)} = [-8.2, -6.5], X_8^{(0)} = [-11.8, -8.0], X_9^{(0)} = [-17.2, -13.5]$$

Test polynomial 5: (Alefeld and Herzberger, 1983)

$$p = \lambda^5 - 30\lambda^4 + 311\lambda^3 - 1278\lambda^2 + 1551\lambda + 630$$

where

$$n = 5, a_1 = 0, a_2 = 3, a_3 = 6, a_4 = 9, a_5 = 12, b_i = 1 (i = 1, \dots, 4)$$

Initial intervals:

$$X_1^{(0)} = [1.9, 3.4], \quad X_2^{(0)} = [4.8, 5.9], \quad X_3^{(0)} = [6.5, 8.1],$$

$$X_4^{(0)} = [8.3, 9.8], \quad X_5^{(0)} = [10.7, 11.9].$$

The tables show the results of all five test polynomials. Table 1 shows the comparison of the number of iteration and CPU time in seconds, between procedures ISS1 and IZSS1-5D.

TABLE 1: Number of Iterations and CPU Times

Polynomial	n	IZSS1		IZSS1-5D	
		No. of iterations	CPU Times (seconds)	No. of iterations	CPU Times (seconds)
1	6	2	0.32813	2	0.23438
2	5	2	0.25000	2	0.21875
3	4	3	0.21875	2	0.15625
4	9	3	0.71875	3	0.703125
5	5	2	0.21875	2	0.171875

TABLE 2: The width of final intervals at iteration k (Polynomial 3, $n = 4$)

Iteration k	Interval width $w_i^{(k)}$	Methods	
		IZSS1	IZSS1-5D
1	$w_1^{(1)}$	0.004440	0.003856
	$w_2^{(1)}$	0.095919	0.035329
	$w_3^{(1)}$	0.002974	0.002906
	$w_4^{(1)}$	0.001150	0.001121
2	$w_1^{(2)}$	3.456×10^{-11}	6.55×10^{-12}
	$w_2^{(2)}$	2.96×10^{-11}	2.13×10^{-12}
	$w_3^{(2)}$	5.00×10^{-14}	5.00×10^{-14}
	$w_4^{(2)}$	1.00×10^{-14}	1.00×10^{-14}
3	$w_1^{(3)}$	1.00×10^{-13}	(converged)
	$w_2^{(3)}$	3.96×10^{-12}	
	$w_3^{(3)}$	5.00×10^{-14}	
	$w_4^{(3)}$	1.00×10^{-14}	

For Table 2, we display the width of intervals at every iteration of the procedures IZSS1-5D and IZSS1 using test polynomial 3, before the algorithm stop at the stopping criterion, $w^{(k)} \leq 10^{-12}$. Table 1 shows that IZSS1-5D converges faster than IZSS1. Note that for test polynomials 3, the procedure IZSS1-5D requires less number of iterations than does procedure IZSS1 except for test polynomials 1, 2, 4 and 5 where the numbers of iterations of both procedures are the same. However, for these four test polynomials, the procedure IZSS1-5D consumes less CPU times compared to the IZSS1 procedure.

5. CONCLUSION

It has been proven that IZSS1-5D has a faster rate of convergence of at least five, whereas the R -order of convergence of IZSS1 is at least four (Rusli *et al.* 2011). Thus, we have this relationship $O_R(\text{IZSS1-5D}, x_i^*) > O_R(\text{IZSS1}, x_i^*)$ ($i=1, \dots, n$). Our proposed procedure IZSS1-5D contributes to shorter CPU times and reduced number of iterations thus enhances the rate of convergence

REFERENCES

- Alefeld, G. and Herzberger, J. (1983). *Introduction to interval computations*, New York: Academic Press.
- Bakar, N. A., Monsi, M. and Hassan, N. (2012). An improved parameter regula falsi method for enclosing a zero of a function. *Applied Mathematical Sciences*. **6**(28): 1347–1361.
- Ehrlic, L. W. (1967). A modified Newton method for polynomials. *Communications of the ACM*. **10**(2): 107-107.
- Jamaludin, N., Monsi, M. and Hassan, N. (2013a). On the interval zero symmetric single step procedure IZSS2-5D for simultaneous bounding of simple polynomial zeros. *International Journal of Mathematical Analysis*. **7**(59): 2941- 2945.
- Jamaludin, N., Monsi, M. and Hassan, N. (2014a). On convergence of the interval zero symmetric single step procedure for polynomial zeros. *International Journal of Mathematical Analysis*. **8**(56): 2769- 2774.

- Jamaludin, N., Monsi, M. and Hassan, N. (2014b). Interval symmetric single-step procedure IZSS2-5D for polynomial zeros. *Sains Malaysiana* **43**(7): 1101- 1104.
- Jamaludin, N., Monsi, M., Hassan, N. and Kartini, S. (2013b). On modified interval symmetric single step procedure ISS2-5D for the simultaneous inclusion of polynomial zeros. *International Journal of Mathematical Analysis*. **7**(20): 983- 988.
- Jamaludin, N., Monsi, M., Hassan, N. and Suleiman, M. (2013c). Modification on interval symmetric single-step procedure ISS-5 δ for bounding polynomial zeros simultaneously. *AIP Conf. Proc.* **1522**: 750-756.
- Jamaludin, N., Monsi, M., Husain, S. K. S. and Hassan, N. (2013d). The interval zero-symmetric single-step procedure IZSS2-5D for the simultaneous bounding of simple polynomial zeros. *AIP Conf. Proc.* **1557**: 268-271.
- Monsi, M., Hassan, N. and Rusli, S. F. (2012). The point zero symmetric single-step procedure for simultaneous estimation of polynomial zeros. *Journal of Applied Mathematics*, Volume 2012, Article ID 709832, 11 pages, doi:10.1155/2012/709832.
- Monsi, M. and Wolfe, M. A. (1988). An algorithm for the simultaneous inclusion of real polynomial zeros. *Applied Mathematics and Computation*. **25**: 333-346.
- Moore, R. E. (1979). *Methods and applications of interval analysis*. Philadelphia: SIAM Publications.
- Ortega, J. M. and Rheinboldt, W. C. (1970). *Iterative solution of nonlinear equations in several variables*. New York: Academic Press.
- Rump, S. M. (1999). *INTLAB-INTERVAL LABORATORY: In Tibor Csendes, Development in reliable computing*. Dordrecht: Kluwer Academic Publisher.

- Rusli, S. F. M., Monsi, M., Hassan, M. A. and Leong, W. J. (2011). On the interval zero symmetric single –step procedure for simultaneous finding of polynomial zeros. *Applied Mathematics Sciences*. **5**: 3693-3706.
- Rusli, S. F. M., Monsi, M. and Hassan, N. (2014a). On the convergence of the Newton's method in the interval symmetric single-step procedure. *International Journal of Mathematical Analysis*. **8**(56): 2775-2779.
- Rusli, S. F. M., Monsi, M. and Hassan, N. (2014b). The Newton's method in the interval symmetric single-step procedure for polynomial zeros. *International Journal of Mathematical Analysis*. **8**(56): 2763-2768.
- Sham, A. W. M., Monsi, M. and Hassan, N. (2013a). An efficient interval symmetric single step procedure ISS1-5D for simultaneous bounding of real polynomial zeros. *International Journal of Mathematical Analysis*. **7**(20): 977-981.
- Sham, A. W. M., Monsi, M. and Hassan, N. (2013b). On the interval zero symmetric single step procedure IZSS1-5D for the simultaneous bounding of real polynomial zeros. *International Journal of Mathematical Analysis*. **7**(59): 2947-2951.
- Sham, A. W. M., Monsi, M., Hassan, N. and Suleiman, M. (2013c). A modified interval symmetric single step procedure ISS-5D. *AIP Conf. Proc.* **1522**: 61–67, doi: 10.1063/1.4801105.